

August 30, 1995

Project No. 301679

Ms. Jean Holdren LITCO P.O. Box 1625, MS 2107 Idaho Falls, ID 83415

Support Documentation: Estimation of Uranium-235 Surface Soil Concentrations

Based on Mass Unrecovered at the BORAX-I Burial Grounds

Dear Jean:

IT is pleased to submit the attached documentation to support estimation of possible uranium (U)-235 surface soil concentrations and activities based on the mass of U-235 unrecovered at the BORAX-I burial ground. This information was developed as a result of public comment on the SL-1 and BORAX-I proposed plan regarding the possibility for soil washing to recover the U-235 remaining at the site. Response to the comment involved estimation of the U-235 concentration in mg/kg. This idea was taken one step further to provide additional information for the uncertainty discussion in Section 6: the concentration of U-235 in mg/kg was converted to pCi/g for comparison to the soil sampling analytical data. The calc briefs detailing the estimates are attached.

Two different soil volumes were used in the estimates to provide a better understanding of the possible range of concentrations at the site. The first volume was based on the entire gravel-covered area (84,000 ft²) and a depth of one foot. The second volume (also one foot deep) was restricted to the areas found with radiological fields above 20 μ R/hr in surveys conducted in 1978 (14,700 ft²). A soil density of 1.5 g/cm³ was assumed, consistent with the density used in the baseline risk assessment for BORAX-I. Results of the calculations are presented in Table 1.

Table 1. Estimated U-235 surface soil concentrations and activities.

Soil Volume (cubic feet)	Concentration (mg/kg)	Activity (pCi/g)	
84,000	1	2	
14,700	6	13	

If you have any questions, please call me at 505-662-1236.

Respectfully submitted,

IT CORPORATION

Richard Filemyr **Project Manager**

attachments

Doug Vetter, IT-Albuquerque Larry Hull, IT-Idaho Falls Central File 301679 CC:

Attachment 1

Calc Brief: BORAX-I Surface Soil Uranium Concentration

BORAX-I Surface Soil Uranium Concentration

Purpose: To determine the concentration of uranium in the surface soils in mg/kg.

Assumptions: The soils contain 3.7 kg of uranium evenly distributed throughout two different

volumes of surface soil: an area of 84,000 ft² to a depth of 1 ft, and 14,700 ft²

to a depth of one foot (see attached figure).

The soil density is 1.5 g/cm³.

Calculation: 84,000 ft3 volume

mg of uranium:

3.7 kg
$$\left(\frac{10^6 \text{ mg}}{\text{kg}}\right)$$
 = 3.7 x 10⁶ mg.

kg of soil:

84,000 ft³
$$\left(\frac{12 \text{ in}}{\text{ft}}\right)^3 \left(\frac{16.387 \text{ cm}^3}{\text{in}^3}\right) = 2,378,605,824 \text{ cm}^3$$

uranium concentration based on 84,000 ft3:

concentration =
$$\frac{3.7 \times 10^6 \text{ mg}}{3,567,908.736 \text{ kg}}$$
 = 1.037 mg/kg

BORAX-I Surface Soil Uranium Concentration Continued

Calculation: 14,700 ft3 volume

kg of soil:

14,700 ft³
$$\left(\frac{12 \text{ in}}{\text{ft}}\right)^3 \left(\frac{16.387 \text{ cm}^3}{\text{in}^3}\right) = 416,256,019.2 \text{ cm}^3$$

1.5 g/cm³
$$(416,256,019.2 \text{ cm}^3) = 624,384,028.8 \text{ g}$$

= 624,384.03 kg


uranium concentration based on 14,700 ft3:

concentration =
$$\frac{3.7 \times 10^6 \text{ mg}}{624,384.03 \text{ kg}}$$
 = 5.9 mg/kg



A,= 32'x66' - 2178 702 151165=9,900 -3= 40 × 40=1,600 74= 25x 16 = 400 = 14,700 ft²
= 14,700 ft²





Assumed gravel boundary

location Approximate security


fence location Grid measurement locations

Measurement locations outside of grid

17 to 20 µR/hr

20 to 50 μP/hr

Gravel covering



Note: The seventh measurement location outside of the grid was not shown in the original documentation.



Exposure rate isoplethic map for the BORAX-I reactor area.

Attachment 2

Calc Brief: BORAX-I Surface Soil Uranium Activity

BORAX-I Surface Soil Uranium Activity

Purpose: To determine the uranium activities in pCi/g based on the concentrations in mg/kg

as determined in Attachment 1.

Reference: Radiological Health Handbook, Revised Edition, January 1970 (see attached).

Given: The specific activity of U-235 is 2.14 x 10⁶ pCi/g (Radiological Health Handbook,

see attached).

Calculation: Based on U-235 soil concentration of 1.037 mg/kg

$$\left(\frac{1.037 \text{ mg}_{(U)}}{\text{kg}_{(\text{soil})}}\right) \left(\frac{1 \text{ g}_{(U)}}{1,000 \text{ mg}_{(U)}}\right) \left(\frac{1 \text{ kg}_{(\text{soil})}}{1,000 \text{ g}_{(\text{soil})}}\right) (2.14 \times 10^8 \text{ pCi/g}_{(U)})$$

= 2.22 pCi/g


Calculation: Based on U-235 soil concentration of 5.9 mg/kg

$$\left(\frac{5.9 \text{ mg}_{(U)}}{\text{kg}_{(\text{soil})}}\right) \left(\frac{1 \text{ g}_{(U)}}{1,000 \text{ mg}_{(U)}}\right) \left(\frac{1 \text{ kg}_{(\text{soil})}}{1,000 \text{ g}_{(\text{soil})}}\right) (2.14 \times 10^6 \text{ pCi/g}_{(U)})$$

= 12.6 pCi/g.

RADIOLOGICAL HEALTH HANDBOOK

Compiled and edited by the Bureau of Radiological Health and the Training Institute Environmental Control Administration



Revised Edition January 1970

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE
Public Health Service
Consumer Protection and Environmental Health Service
Rockville, Maryland 20852

For said by the Superintendent of Donamonia, T.L. Government Printing Office Washington, D.C. 2005 - Price 84.00

SPECIFIC ACTIVITY

Radionuclida	Half-Life	Curies per gram	Radionuclid.	Half-Life	Curies per gram
Hydrogen-3	12.3y	9.54×10 ³	Molybdenum-99	67h	4.72×10 ⁵
Carbon-14	5730y	4.46	Tachnetium-99m	6.Oh	5.28×10°
Nitrogen-16	7.2.	9.79×10 ⁻⁰	Ruthenium-106	367d	3.36x10 ³
Sodium-22	2.60y	6.25×10 ³	Iodine-125	60 d	1.74x10 ⁴
Sodium-24	15.0h	8.71x10 ⁶	Iodine-130	12.4h	1.94×10 ⁶
Phosphorus-32	14.34	21.85×10 ⁵	Iodine-131	8.054	1.24×10 ⁵
Sulfur-35	884	4.24x10 ⁴	Barium-133	7.2y	374
Chlorine-36	3.1x10 ⁸ y	3.21x10 ⁻⁸	Cesium-134	2.05y	1.30×10 ^a
Argon-41	1.83h	4.18x10 ⁷	Cesium-137	30.0y	87.0
Potessium-42	12.4h	6.02x10 ⁶	Barium-140	12.8d	7.29×10 ⁴
Calcium-45	1654	1.76x10 ⁴	Lanthanum-140	40.22h	5.57×10 ⁵
Chromium-51	27.8d	9.21x10 ⁴	Carium-141	33 d	2.81×10 ⁴
Manganese-54	3034	7.98×10 ³	Carium-144	284d	3.19×10 ³
Iron-55	2.6y·	2.50×10 ²	Praseodymium-144	17.3m	7.55×10 ⁷
Manganesa-56	2.576h	2,17×10 ⁷	Promethium-147	2.62 y	929
Cobalt-57	2704	8.48×10 ³	Tantalum-182	115d	6.24×10 ³
Iron-59	45d	4,92×10	Tungsten-185	75 4	9,41x10 ³
Nickel-59	8×10 ⁶ y	7.58×10 ⁻²	Iridium-192	74.2d	9.17x10 ³
Cobalt-60	5.26y	1.13×10 ³	Gold-198	64.8h	2.44×10 ⁵
Nickel-63	92y	61.7	Gold-199	·75.6h	2.08×10 ⁵
Copper-64	12.8h	3.83×10 ⁸	Marcury-203	46.9d	1.37×10 ⁴
Zinc-65	2454	8.20×10 ³	Thellium-204	3.8y	462
Gallium-72	14.1h	3.09×10 ⁶	Polonium-210	138.4d	4.49x10 ³
Arsenic-76	26.5h	1.56×10 ⁶	Poionium-212	304ns	1,75×10 ¹³
Bromine-82	35.34h	1.08×10 ⁸	Radium-226	1602y	0.988
Rubidium-86	18.66d	8.14×10 ⁴	Thorium-232	1.41x10 ¹⁰ y	1.09×10
Strontium-89	524	2.82×10 ⁴	.Uranium-233	1.62x10 ⁸ y	9.48x10 ⁻¹
Strontium-90	28.1y	141	Thorium-234	24.1d	2.32×10 ⁴
Yttrium-90	64h	5.44×10	Uranium-23/5	7.1x10 ⁴ y	2.14×10
Ycerium-91	58.84	2.44×10 ⁴	Uranium-238	4.51x10 ⁸ y	3.33×10
			Plutonium-239	2.44x10 ⁴ y	6.13x10°